


Cough (acute): antimicrobial prescribing



Cough (acute): antimicrobial prescribing

Choice of antibiotic: adults aged 18 years and over

| Antibiotic ¹ | Dosage and course length ² |
|---|--|
| First choice | |
| Doxycycline ³ | 200 mg on first day, then 100 mg once a day for 4 days (5-day course in total) |
| Alternative first choices ⁴ | |
| Amoxicillin | 500 mg three times a day for 5 days |
| Clarithromycin | 250 mg to 500 mg twice a day for 5 days |
| Erythromycin | 250 mg to 500 mg four times a day or 500 mg to 1000 mg twice a day for 5 days |
| ¹ See BNF for appropriate use and dosing in specific populations, for example, hepatic impairment, renal impairment, pregnancy and breast-feeding ² Doses given are by mouth using immediate-release medicines, unless otherwise stated ³ Doxycycline should not be given to pregnant women, and the possibility of pregnancy should be considered in women of childbearing age (BNF , December 2018) ⁴ Amoxicillin or erythromycin are preferred in women who are pregnant | |

When exercising their judgement, professionals and practitioners are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or the people using their service. It is not mandatory to apply the recommendations, and the guideline does not override the responsibility to make decisions appropriate to the circumstances of the individual, in consultation with them and their families and carers or guardian.

Choice of antibiotic: children and young people under 18 years

| Antibiotic ¹ | Dosage and course length ² |
|--|--|
| First choice | |
| Amoxicillin | 1 to 11 months: 125 mg three times a day for 5 days 1 to 4 years: 250 mg three times a day for 5 days 5 to 17 years: 500 mg three times a day for 5 days |
| Alternative first choices ³ | |
| Clarithromycin | 1 month to 11 years: Under 8 kg, 7.5 mg/kg twice a day for 5 days 8 to 11 kg, 62.5 mg twice a day for 5 days 12 to 19 kg, 125 mg twice a day for 5 days 20 to 29 kg, 187.5 mg twice a day for 5 days 30 to 40 kg, 250 mg twice a day for 5 days 12 to 17 years: 250 mg to 500 mg twice a day for 5 days |
| Erythromycin | 1 month to 1 year: 125 mg four times a day or 250 mg twice a day for 5 days 2 to 7 years: 250 mg four times a day or 500 mg twice a day for 5 days 8 to 17 years: 250 mg to 500 mg four times a day or 500 mg to 1000 mg twice a day for 5 days |
| Doxycycline ⁴ | 12 to 17 years: 200 mg on first day, then 100 mg once a day for 4 days (5-day course in total) |
| ¹ See BNF for children for appropriate use and dosing in specific populations, for example, hepatic impairment and renal impairment ² The age bands apply to children of average size and, in practice, the prescriber will use the age bands in conjunction with other factors such as the severity of the condition and the child's size in relation to the average size of children of the same age. Doses given are by mouth using immediate-release medicines, unless otherwise stated ³ Amoxicillin or erythromycin are preferred in young women who are pregnant ⁴ Doxycycline should not be given to young women who are pregnant, and the possibility of pregnancy should be considered in young women of childbearing age (BNF for children , December 2018) | |