

PENNINE LANCASHIRE OPIOID CONVERSION TABLE - A GUIDE TO EQUIVALENT DOSES OF OPIOID DRUGS


The table below has been adapted from the Lancashire and South Cumbria Clinical Practice Summary Nov 2021 for local use within Pennine Lancashire, to include dose ranges and to round doses to ease administration.

Use the table as a guide (not a set of definitive equivalences) to identify an appropriate starting point for your prescribing decisions. **ALL** prescribing decisions must be based on a **full clinical assessment**. **Higher opioid doses may be needed for some patients – seek specialist palliative care advice.**

Considering **reducing prescribed opioid dose by 30-50%** if converting from one route to another route (e.g. transdermal to oral to oral to subcutaneous) or if there is concern about **opioid toxicity** (confusion, drowsiness, myoclonic jerks, slowed respiration, pin-point pupils). See Lancashire and South Cumbria Clinical Practice Summary Nov 2021 for guidance on conversion to a transdermal fentanyl patch or CSCI (syringe pump). Consider the role of adjuvant medication before rotating opioids, changing the dose or route.

Never increase an opioid dose by more than 50% of the previous 24 hour regular dose without SPECIALIST ADVICE.

Consider prescribed doses of moderate opioids (Codeine and Tramadol). Factor these in when converting to regular morphine (or another strong opioid) or when calculating PRN dosages.



| Route | Morphine (mg) | | | | | Oxycodone (mg) | | | | | Fentanyl Patch (mcg/hr) | Buprenorphine Patch (mcg/hr) | Alfentanil Syringe pump over 24hrs (not used PRN SC, consider oxycodone PRN) |
|--|--|----------------|---------|-------------------------|----------|----------------|----------------|----------|-------------------------|----------------|-------------------------|------------------------------|--|
| | Oral | | | SC | | Oral | | | SC | | | | |
| | 24hr Total | 12hrly MR Dose | PRN | Syringe Pump over 24hrs | PRN | 24hr Total | 12hrly MR Dose | PRN | Syringe Pump over 24hrs | PRN | | | |
| Dose | 20 | 10 | 2.5 - 5 | 10 | 2.5 | 10 | 5 | 2.5 | 5 | 1 | | | 500micrograms |
| | 30 | 15 | 5 | 15 | 2.5 | 15 | * | 2.5 | 7.5 | 1 - 2 | 12 micrograms | 10 micrograms | |
| | 40 | 20 | 5 - 10 | 20 | 2.5 - 5 | 20 | 10 | 2.5 - 5 | 10 | 2 | - | - | 1mg |
| | 50 | 25 | 5 - 10 | 25 | 2.5 - 5 | 25 | * | 2.5 - 5 | 10 - 15 | 2.5 | - | 20 micrograms | |
| | 60 | 30 | 10 | 30 | 5 | 30 | 15 | 5 | 15 | 2.5 | 25 micrograms | - | 1.5mg |
| | 70 | 35 | 10 - 15 | 35 | 5 - 7.5 | 35 | * | 5 - 7.5 | 20 | 2.5 - 5 | - | 30 micrograms | |
| | 80 | 40 | 10 - 15 | 40 | 5 - 7.5 | 40 | 20 | 5 - 7.5 | 20 | 2.5 - 5 | - | 35 micrograms | 2mg |
| | 100 | 50 | 15 - 20 | 50 | 7.5 - 10 | 50 | 25 | 7.5 - 10 | 25 | 2.5 - 5 | - | - | 2.5mg |
| | 120 | 60 | 20 | 60 | 10 | 60 | 30 | 10 | 30 | 5 | 50 micrograms | 52.5 micrograms | 3mg |
| | Seek Specialist advice for higher doses | | | | | | | | | | | | |
| 160 | 80 | 25 - 30 | 80 | 10 - 15 | 80 | 40 | 10-15 | 40 | 5 - 10 | 75 micrograms | 70 micrograms | 4mg | |
| 240 | 120 | 40 | 120 | 20 | 120 | 60 | 20 | 60 | 10 | 100 micrograms | - | 6mg | |
| * When equal divided doses not possible due to tablet strength e.g. Oxycodone 25mg/24hrs. Prescribe doses at higher or lower level e.g. 10mg BD or 15mg BD, dependant on clinical judgement* | | | | | | | | | | | | | |